arXiv:1509.06399v1 [math.CA] 21 Sep 2015

A NOTE ON A MULTILINEAR LOCAL Tb THEOREM FOR CALDERÓN-ZYGMUND OPERATORS

JAROD HART AND LUCAS OLIVEIRA

ABSTRACT. In this short note, we extend a local Tb theorem that was proved in [3] to a full multilinear local Tb theorem.

1. INTRODUCTION

In [3], we proved a multilinear local Tb theorem for square functions, and applied it to prove a local Tb theorem for singular integrals. The local Tb theorem for singular integrals had "local Tb type" testing conditions for the operator T on pseudo-accretive collections $\{b_Q^i\}$ for $i = 1,\dots,m$, but tested the adjoints of T on the constant function 1; see [3] for more details on this. In this note, we make a few observations to show that this result can be extended to the following local Tb theorem for multilinear singular integral operators.

Theorem 1.1. *Let T be a continuous m linear operator from $\mathscr{S}\times\cdots\times\mathscr{S}$ into $\mathscr{S}'$ with a standard Calderón-Zygmund kernel K. Suppose that $T \in WBP$ and there exist $2 \le q < \infty$ and $1 < q_{i,j} < \infty$ with $\frac{1}{q_j} = \sum_{i=1}^m \frac{1}{q_{i,j}}$ and an m-compatible collection of functions $\{b_Q^{i,j}\}$ indexed by dyadic cubes Q and $j = 1,\dots,m$ such that*

$$\int_Q \left(\int_0^{\ell(Q)} |Q_t T^{*j}(P_t b_Q^{1,j},\dots,P_t b_Q^{m,j})(x)|^2 \frac{dt}{t} \right)^{\frac{q_j}{2}} dx \lesssim |Q|. \tag{1}$$

Then T is bounded from $L^{p_1}\times\cdots\times L^{p_m}$ into L^p for all $1 < p_i < \infty$ such that $\frac{1}{p} = \frac{1}{p_1} + \cdot + \frac{1}{p_m}$. Here we assume that this estimate holds for any approximation to the identity P_t with smooth compactly supported kernels and any Littlewood-Paley-Stein projection operators Q_t whose kernels also are smooth compactly supported function.

The precise meaning of (1) is the following: For any $\varphi,\psi \in C_0^\infty$ such that $\widehat{\varphi}(0) = 1$ and $\widehat{\psi}(0) = 0$, (1) holds for $P_t f = \varphi_t * f$, $Q_t f = \psi_t * f$, $\varphi_t(x) = \frac{1}{t^n}\varphi(\frac{x}{t})$, and $\psi_t(x) = \frac{1}{t^n}\psi(\frac{x}{t})$, where the constant is independent of the dyadic cube Q, but may depend on φ and ψ.

We prove this theorem by applying the square function estimates that were also proved in [3], but with a few minor modifications to allow for the extension to Theorem 1.1. This note is intended to be an addendum to the article [3]. So the reader should refer to that article for definitions, discussion, and history related to these results.

Date: September 23, 2015.
2010 *Mathematics Subject Classification.* 42B20, 42B25, 42B30.
Key words and phrases. Square Function, Littlewood-Paley-Stein, Calderón-Zygmund Operators.

2. A Few Definitions and Results from [3]

Define the family of multilinear of operators $\{\Theta_t\}_{t>0}$

$$\Theta_t(f_1,\dots,f_m)(x)=\int_{\mathbb{R}^{mn}}\theta_t(x,y_1,\dots,y_m)\prod_{i=1}^m f_i(y_i)dy_i \tag{2}$$

where $\theta_t:\mathbb{R}^{(m+1)n}\to\mathbb{C}$ and the square function

$$S(f_1,\dots,f_m)(x)=\left(\int_0^\infty|\Theta_t(f_1,\dots,f_m)(x)|^2\frac{dt}{t}\right)^{\frac{1}{2}} \tag{3}$$

associated to $\{\Theta_t\}_{t>0}$, where f_i for $i=1,\dots,m$ are initially functions in $C_0^\infty(\mathbb{R}^n)$. Also assume that θ_t satisfies for all $x,y_1,\dots,y_m,x',y_1',\dots,y_m'\in\mathbb{R}^n$

$$|\theta_t(x,y_1,\dots,y_m)|\lesssim\frac{t^{-mn}}{\prod_{i=1}^m(1+t^{-1}|x-y_i|)^{N+\gamma}} \tag{4}$$

$$|\theta_t(x,y_1,\dots,y_m)-\theta_t(x,y_1,\dots,y_i',\dots,y_m)|\lesssim t^{-mn}(t^{-1}|y_i-y_i'|)^\gamma \tag{5}$$

$$|\theta_t(x,y_1,\dots,y_m)-\theta_t(x',y_1,\dots,y_m)|\lesssim t^{-mn}(t^{-1}|x-x'|)^\gamma \tag{6}$$

for some $N>n$ and $0<\gamma\le 1$. The following results was proved in [3].

Theorem. *Let* Θ_t *and* S *be defined as in* (2) *and* (3) *where* θ_t *satisfies* (4)-(6). *Suppose there exist* $q_i,q>1$ *for* $i=1,\dots,m$ *with* $\frac{1}{q}=\sum_{i=1}^m\frac{1}{q_i}$ *and functions* b_Q^i *indexed by dyadic cubes* $Q\subset\mathbb{R}^n$ *for* $i=1,\dots,m$ *such that for every dyadic cube* Q

$$\int_{\mathbb{R}^n}|b_Q^i|^{q_i}\le B_1|Q| \tag{7}$$

$$\frac{1}{B_2}\le\left|\frac{1}{|Q|}\int_Q\prod_{i=1}^m b_Q^i(x)dx\right| \tag{8}$$

$$\left|\frac{1}{|R|}\int_R\prod_{i=1}^m b_Q^i(x)dx\right|\le B_3\prod_{i=1}^m\left|\frac{1}{|R|}\int_R b_Q^i(x)dx\right| \quad \textit{for all dyadic subcubes } R\subset Q \tag{9}$$

$$\int_Q\left(\int_0^{\ell(Q)}|\Theta_t(b_Q^1,\dots,b_Q^m)(x)|^2\frac{dt}{t}\right)^{\frac{q}{2}}dx\le B_4|Q|. \tag{10}$$

Then S *satisfies*

$$||S(f_1,\dots,f_m)||_{L^p}\lesssim\prod_{i=1}^m||f||_{L^{p_i}} \tag{11}$$

for all $1<p_i<\infty$ *and* $2\le p<\infty$ *satisfying* $\frac{1}{p}=\frac{1}{p_1}+\cdots+\frac{1}{p_m}$.

If $\{b_Q\}$ satisfies (7) and (8), we say that $\{b_Q\}$ is a pseudo-accretive system. We say that $\{b_Q^i\}$ for $i=1,\dots,m$ is an m-compatible, or just compatible, collection of pseudo-accretive systems if they satisfy (7)-(9).

3. PROOF OF THEOREM 1.1

Proof of Theorem 1.1. Let P_t be a smooth approximation to identity operators with smooth compactly supported kernels that $P_t^2 f \to f$ as $t \to 0^+$ and $P_t^2 f \to 0$ as $t \to \infty$ in $\mathscr{S}$ for $f \in \mathscr{S}_0$. Here $\mathscr{S}_0$ is the subspace of Schwartz functions satisfying $|\widehat{f}(\xi)| \le C_M|\xi|^M$ for all $M \in \mathbb{N}$. There exist Littlewood-Paley-Stein projection operators $Q_t^{(i)}$ for $i = 1,2$ with smooth compactly supported kernels such that $t\frac{d}{dt}P_t^2 = Q_t^{(2)}Q_t^{(1)} = Q_t$. Using these operators, we decompose T for $f_i \in \mathscr{S}_0$, $i = 0,\dots,m$

$$
\begin{aligned}
\langle T(f_1,\dots,f_m), f_0\rangle &= \int_0^\infty t\frac{d}{dt}\left\langle T(P_t^2 f_1,\dots,P_t^2 f_m), P_t^2 f_0\right\rangle \frac{dt}{t}\\
&= \sum_{i=0}^m \left\langle \int_0^\infty Q_t T(P_t^2 f_1,\dots,P_t f_{i-1},P_t f_0,P_t f_{i+1}\dots,P_t^2 f_m)\frac{dt}{t}, f_i\right\rangle\\
&= \sum_{i=0}^m \langle T_i(f_1,\dots,f_{i-1},f_0,f_{i+1},\dots,f_m), f_i\rangle, \qquad (12)
\end{aligned}
$$

where we take the last line in (12) as the definition of T_i for $i = 0,1,\dots,m$. It follows that T_i is a multilinear singular integral operator with standard kernel

$$K_i(x,y_1,\dots,y_m) = \int_0^\infty \left\langle T^{*i}(\varphi_t^{y_1},\dots,\varphi_t^{y_m}),\psi_t^x\right\rangle \frac{dt}{t},$$

Also let Θ_t^i be the multilinear operator associated to

$$\theta_t^i(x,y_1,\dots,y_m) = Q_t^{(1)}T^{*i}(\varphi_t^{y_1},\dots,\varphi_t^{y_m})(x),$$

and let S_i be the square function associated to Θ_t^i. Note that $\theta_t^i(x,y_1,\dots,y_m) \ne \left\langle T^{*i}(\varphi_t^{y_1},\dots,\varphi_t^{y_m}),\psi_t^x\right\rangle$, and that T_i is not actually the integral of Θ_t^i (one has Q_t and the other has $Q_t^{(1)}$). Furthermore, by the hypotheses on T^{*i} and by the local Tb theorem for square functions from [3], it follows that S_i is bounded from $L^{2m}\times\cdots\times L^{2m}$ into L^2. Therefore we have

$$
\begin{aligned}
\langle T_i(f_1,\dots,f_m), f_0\rangle &= \int_0^\infty \int_{\mathbb{R}^n} Q_t T(P_t^2 f_1,\dots,P_t^2 f_m)(x) f_0(x)dx\frac{dt}{t}\\
&= \int_0^\infty \int_{\mathbb{R}^n} Q_t^{(1)} T(P_t^2 f_1,\dots,P_t^2 f_m)(x) Q_t^{(2)*} f_0(x)dx\frac{dt}{t}\\
&\le ||S_i(f_1,\dots,f_m)||_{L^2}\left|\left|\left(\int_0^\infty |Q_t^{(2)*}f_0|^2\frac{dt}{t}\right)^{\frac{1}{2}}\right|\right|_{L^2} \lesssim ||f_1||_{L^{2m}}\cdots||f_m||_{L^{2m}}||f_0||_{L^2}.
\end{aligned}
$$

Hence T_i is bounded from $L^{2m}\times\cdots\times L^{2m}$ into L^2, and by the multilinear Calderón-Zygmund theory developed by Grafakos and Torres in [1, 2], it follows that T_i also bounded from $L^{p_1}\times\cdots\times L^{p_m}$ into L^p for all $1 < p_1,\dots,p_m < \infty$ with $\frac{1}{p} = \frac{1}{p_1}+\cdots+\frac{1}{p_m}$. In particular, it follows that T_i is bounded from $L^{m+1}\times\cdots\times L^{m+1}$ into $L^{\frac{m+1}{m}}$ for each $i=0,1,\dots,m$. Then continuing from (12), we have

$$\begin{aligned}|\langle T(f_1,\dots,f_m),f_0\rangle| &\le \sum_{i=0}^{m}|\langle T_i(f_1,\dots,f_{i-1},f_0,f_{i+1},\dots,f_m),f_i\rangle|\\ &\le \sum_{i=0}^{m}||T_i(f_1,\dots,f_{i-1},f_0,f_{i+1},\dots,f_m)||_{L^{\frac{m+1}{m}}}||f_i||_{L^{m+1}}\lesssim\prod_{j=0}^{m}||f_j||_{L^{m+1}}.\end{aligned}$$

Therefore T is bounded from $L^{m+1}\times\cdots L^{m+1}$ into $L^{\frac{m+1}{m}}$. Then it again follows from the multilinear Calderón-Zygmund theory in [1, 2] that T is bounded from $L^{p_1}\times\cdots\times L^{p_m}$ into L^p for all $1<p_1,\dots,p_m<\infty$ such that $\frac{1}{p}=\frac{1}{p_1}+\cdots+\frac{1}{p_m}$. □

REFERENCES

1. L. Grafakos, R.H. Torres, *On multilinear singular integrals of Calderón-Zygmund type*, Publ. Mat., (2002) 57-91.
2. L. Grafakos and R. H. Torres, *Multilinear Calderón–Zygmund theory*, Adv. in Math. **165** (2002), 124–164.
3. A. Grau de la Herrán, J. Hart, and L. Oliveira, *Multilinear local Tb thoerem for square functions*, Annales Academiæ Scientiarum Fennicæ Mathematica, **38**, (2013), 1-24.

HIGUCHI BIOSCIENCES CENTER, UNIVERSITY OF KANSAS, LAWRENCE, KS
E-mail address: `jvhart@ku.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL, PORTO ALEGRE, RIO GRANDE DO SUL, BRAZIL
E-mail address: `lucas.oliveira@ufrgs.br`